

CS4311

Design and Analysis of Algorithms

Tutorial for Fun:
Deriving Catalan Number Formula

Generating Function

- Let $S = s_0, s_1, s_2, \dots$ be a series of numbers we are interested
- Then the function

$$F(x) = \sum s_i x^i = s_0 + s_1 x + s_2 x^2 + s_3 x^3 + \dots$$

is called a **generating function** of S

Generating Function

Example 1:

$$F(x) = \sum i x^i = x + 2x^2 + 3x^3 + \dots$$

is the generating function of $0, 1, 2, \dots$

Example 2:

$$F(x) = 1 + 4x + 6x^2 + 4x^3 + x^4$$

is the generating function of

$$\binom{4}{0}, \binom{4}{1}, \binom{4}{2}, \binom{4}{3}, \binom{4}{4}$$

Closed Form

- Sometimes, generating function can be expressed in the **closed form** :

Example 1:

$$F(x) = \sum x^i = 1 + x + x^2 + x^3 + \dots$$

has a closed form $1 / (1-x)$

Why? Because $(1-x)(1 + x + x^2 + x^3 + \dots) = 1$

Closed Form

Example 2:

$$\begin{aligned} F(x) &= \sum C(n,i) x^i \\ &= 1 + nx + C(n,2)x^2 + \dots nx^{n-1} + x^n \end{aligned}$$

has a closed form $(1+x)^n$

Example 3: How about the closed form of

$$F(x) = \sum i x^i = x + 2x^2 + 3x^3 + \dots ?$$

Closed Form

- Generating function is very useful in
(I) solving combinatorial problems, and
(II) solving recurrences
- Usually, the closed form is important because it can simplify the notation a lot!
- We will see how generating function is used to get Catalan number formula

Catalan Number

- Let us define the n^{th} Catalan number
 - $c_n = \#$ binary trees with n internal nodes
 - $= \#$ binary trees with $n+1$ leaves
- What is c_0, c_1, c_2, c_3 ?


$$c_0 = 1$$




$$c_1 = 1$$



$$c_2 = 2$$



$$c_3 = 5$$



Catalan Number

- Note: an n -node tree can be formed by:
 - (i) choosing the k^{th} node to be its root
 - (ii) arrange the left tree in any order
 - (iii) arrange the right tree in any orderSo, there are $C_{k-1} * C_{n-k}$ choices

$$\begin{aligned} \rightarrow C_n &= C_0 C_{n-1} + C_1 C_{n-2} + C_2 C_{n-3} + \dots + C_{n-1} C_0 \\ &= \sum_{k=1 \text{ to } n} C_{k-1} C_{n-k} \end{aligned}$$

Catalan Number

So, we have:

$$C_0 = 1$$

$$C_1 = C_0 C_0 x$$

⋮

$$C_{n-1} x^{n-1} = \sum_{k=1 \text{ to } n-1} C_{k-1} C_{n-k} x^{n-1}$$

$$C_n x^n = \sum_{k=1 \text{ to } n} C_{k-1} C_{n-k} x^n$$

⋮

Catalan Number

$$\begin{aligned}\text{Let } F(x) &= \text{generating function of Catalan \#} \\ &= c_0 + c_1x + \dots + c_nx^n + \dots \\ &= \text{sum of LHS}\end{aligned}$$

However,

sum of RHS

$$\begin{aligned}&= 1 + x \left[c_0c_0 + (c_0c_1 + c_1c_0)x + \dots \right. \\ &\quad \left. + (c_0c_{n-1} + \dots + c_{n-1}c_0)x^{n-1} + \dots \right] \\ &= 1 + x (F(x))^2\end{aligned}$$

Catalan Number

Thus,

$$F(x) = 1 + x (F(x))^2$$

Or, $x (F(x))^2 - F(x) + 1 = 0$

Hence, we get a closed form of $F(x)$:

$$\begin{aligned} F(x) &= (1 \pm \sqrt{1 - 4x}) / (2x) \\ &= (1 - \sqrt{1 - 4x}) / (2x) \quad (\text{why?}) \end{aligned}$$

Catalan Number

$$\text{Let } C\left(\frac{1}{2}, k\right) = \frac{1}{2} \left(\frac{1}{2} - 1\right) \left(\frac{1}{2} - 2\right) \dots \left(\frac{1}{2} - k + 1\right) / k!$$

Then, by binomial expansion, (or Taylor)

$$(1 - 4x)^{1/2}$$

$$= 1 + \frac{1}{2}(-4x) + \dots + C\left(\frac{1}{2}, n\right)(-4x)^n + \dots$$

$$= 1 - 2x - \dots - 4^n \frac{1}{2} \left(1 - \frac{1}{2}\right) \left(2 - \frac{1}{2}\right) \dots \left(n - 1 - \frac{1}{2}\right) x^n / n!$$

- ...

Simplifying Terms

We claim that:

$$4^n \frac{1}{2} \left(1 - \frac{1}{2}\right) \left(2 - \frac{1}{2}\right) \dots \left(n - 1 - \frac{1}{2}\right) / n! \\ = C(2n, n) / (2n-1)$$

$$\rightarrow (1-4x)^{1/2} = 1 - 2x - \dots - C(2n, n)x^n / (2n-1) - \dots$$

$$\rightarrow F(x) = 1 - ((1-4x)^{1/2}) / (2x) \\ = 1 + \dots + C(2n, n)x^{n-1} / (2(2n-1)) + \dots$$

Simplifying Terms

Proof of claim:

$$\begin{aligned} & 4^n \frac{1}{2} \left(1 - \frac{1}{2}\right) \left(2 - \frac{1}{2}\right) \dots \left(n - 1 - \frac{1}{2}\right) / n! \\ &= 2^n (1)(1)(3)(5)\dots(2n-3) / n! \\ &= 2^n n! (1)(3)(5)\dots(2n-3)(2n-1) / (n! n!(2n-1)) \\ &= (2)(4)(6)\dots(2n)(1)(3)(5)\dots(2n-1) / (n! n!(2n-1)) \\ &= (2n)! / (n! n! (2n-1)) \end{aligned}$$

Catalan Number

Recall: n^{th} Catalan number c_n
= coefficient of x^n in $F(x)$

$$\begin{aligned}\rightarrow c_n &= C(2n+2, n+1) / (2(2n+1)) \\ &= (2n+2)! / ((n+1)! (n+1)! 2(2n+1)) \\ &= (2n+2)! / (n! (n+1)! (2n+2)(2n+1)) \\ &= (2n)! / (n! (n+1)!) \\ &= (2n)! / (n! n! (n+1)) \\ &= C(2n, n) / (n+1)\end{aligned}$$